

Development of a Multimarker Assay for Early Detection of Ovarian Cancer

Zoya Yurkovetsky, Steven Skates, Aleksey Lomakin, Brian Nolen, Trenton Pulsipher, Francesmary Modugno, Jeffrey Marks, Andrew Godwin, Elieser Gorelik, Ian Jacobs, Usha Menon, Karen Lu, Donna Badgwell, Robert C. Bast Jr, and Anna E. Lokshin

See accompanying editorial on page 2128

ABSTRACT

Purpose

Early detection of ovarian cancer has great promise to improve clinical outcome.

Patients and Methods

Ninety-six serum biomarkers were analyzed in sera from healthy women and from patients with ovarian cancer, benign pelvic tumors, and breast, colorectal, and lung cancers, using multiplex xMAP bead-based immunoassays. A Metropolis algorithm with Monte Carlo simulation (MMC) was used for analysis of the data.

Results

A training set, including sera from 139 patients with early-stage ovarian cancer, 149 patients with late-stage ovarian cancer, and 1,102 healthy women, was analyzed with MMC algorithm and cross validation to identify an optimal biomarker panel discriminating early-stage cancer from healthy controls. The four-biomarker panel providing the highest diagnostic power of 86% sensitivity (SN) for early-stage and 93% SN for late-stage ovarian cancer at 98% specificity (SP) was comprised of CA-125, HE4, CEA, and VCAM-1. This model was applied to an independent blinded validation set consisting of sera from 44 patients with early-stage ovarian cancer, 124 patients with late-stage ovarian cancer, and 929 healthy women, providing unbiased estimates of 86% SN for stage I and II and 95% SN for stage III and IV disease at 98% SP. This panel was selective for ovarian cancer showing SN of 33% for benign pelvic disease, SN of 6% for breast cancer, SN of 0% for colorectal cancer, and SN of 36% for lung cancer.

Conclusion

A panel of CA-125, HE4, CEA, and VCAM-1, after additional validation, could serve as an initial stage in a screening strategy for epithelial ovarian cancer.

J Clin Oncol 28:2159-2166. © 2010 by American Society of Clinical Oncology

INTRODUCTION

Ovarian cancer is the fourth most frequent cause of death from cancer in women in Europe and the United States.¹⁻³ Because ovarian cancers typically cause few specific symptoms, more than 70% of patients are diagnosed with advanced disease, where 5-year survival rates are less than 30%.^{1,3} In contrast, the 25% of patients who are diagnosed with stage I disease have a 5-year survival rate of up to 90%, and patients with stage II disease have a 5-year survival rate of up to 70%.^{2,3} Therefore, early detection of ovarian cancer has great promise to improve clinical outcome.

At present, no screening techniques are recommended for early detection of ovarian cancer in the general population. CA-125, the most frequently

used serum biomarker for ovarian cancer, has a sensitivity (SN) of only 50% to 60% for early-stage disease in postmenopausal women when specificity (SP) is set at 99%.⁴⁻⁶ Transvaginal sonography (TVS), computed tomography, magnetic resonance imaging, and power Doppler offer less than 90% SN for early ovarian cancer, and their expense and relatively high false-positive rates preclude annual screening.⁷⁻⁹ Considering the low prevalence of ovarian cancer, a screening strategy must achieve a minimum SP of 99.6% and an SN of more than 75% for early-stage disease to avoid an unacceptable level of false-positive results and achieve a positive predictive value of 10%.^{10,11} Using TVS as a second-line test, previous CA-125-based screening studies indicate that a first-line SP of 98% for an annual test could assure required SP (> 99.6%) and positive

From the University of Pittsburgh Cancer Institute; Departments of Epidemiology, Pathology, Medicine, and Obstetrics/Gynecology and Reproductive Sciences, University of Pittsburgh, Pittsburgh; Fox Chase Cancer Center, Philadelphia, PA; Biostatistics Center, Massachusetts General Hospital; Harvard Medical School, Boston; Department of Physics, Massachusetts Institute of Technology, Cambridge, MA; Department of Surgery, Duke University Medical Center, Durham, NC; Departments of Gynecologic Oncology and Experimental Therapeutics, The University of Texas M. D. Anderson Cancer Center, Houston, TX; and Department of Gynecological Oncology, Institute of Women's Health, University College London, London, United Kingdom.

Submitted August 3, 2008; accepted January 5, 2010; published online ahead of print at www.jco.org on April 5, 2010.

Supported by an Early Detection Research Network Associate Member Award and Grants No. R01 CA098642, R01 CA108990, P50 CA083639, CA086381, and CA105009 from the National Institutes of Health NIH; Avon (NIH/National Cancer Institute); University of Pittsburgh Cancer Institute Hillman Fellows Award; and The Frieda G. and Saul F. Shapira BRCA Cancer Research Program Award (A.E.L.).

Authors' disclosures of potential conflicts of interest and author contributions are found at the end of this article.

Corresponding author: Anna E. Lokshin, MD, University of Pittsburgh Hillman Cancer Center, 5117 Centre Ave, Pittsburgh, PA 15213; e-mail: lokshina@pitt.edu.

© 2010 by American Society of Clinical Oncology

0732-183X/10/2813-2159/\$20.00

DOI: 10.1200/JCO.2008.19.2484

Table 1. Characteristics of the Patient Population

Histology	Training Set 2											Validation Set												
	No. of Patients	Race (No. of patients)					Grade (No. of patients)			Age (years)			No. of Patients	Race (No. of patients)					Grade (No. of patients)			Age (years)		
		Asian	Black	Hispanic	White	1	2	3	Range	Median	Mean	Asian		Black	Hispanic	White	1	2	3	Range	Median	Mean		
Healthy postmenopausal	1,102	52	87	42	921				48-87	56	57.8	929	42	75	23	789				48-77	55	55.4		
Ovarian cancer, postmenopausal, stages IA-IIB	139	5	6	126	2	53	35	51	45-85	59	61.2	44	2	3	1	38	5	10	29	47-87	62	62.6		
Endometrioid	42											20												
Mucinous	29											5												
Serous	27											10												
Clear cell	20											8												
Other	21											1												
Ovarian cancer postmenopausal stages IIC-IV	149					42	50	57	48-87	66	65.1	124	3	4	3	95	2	47	75	48-87	64	65.6		
Endometrioid												39												
Mucinous												4												
Serous												47												
Clear cell												11												
Other												5												
Benign postmenopausal												296	12	16	9	259				49-85	64	62.7		
Breast cancer												210	7	50	6	147				53-82	63	64.4		
Lung cancer												74	1	4	2	67				47-91	69	67.8		
Colorectal cancer												31	0	2	1	28				49-86	63	64.0		

predictive value ($> 10\%$) and would reduce the number of ultrasound examinations performed annually to a cost-effective level of 2%.^{10,11}

Similar to CA-125, several other individual ovarian cancer-associated serum protein biomarkers lack sufficient SN or SP for detection of early-stage disease.¹²⁻¹⁶ Recently, combinations of serum tumor markers have achieved greater SN than individual markers, while maintaining high SP. Two combinations, CA-125, CA 72-4, CA 15-3, and M-CSF¹⁷ and CA-125, apolipoprotein A1, truncated form of transthyretin, and a cleavage fragment of inter- α -trypsin inhibitor heavy chain H4,¹⁸ substantially improved test accuracy over CA-125 alone, with SNs of 70% to 73% at an SP of 97% to 98%. A panel of six biomarkers (CA-125, leptin, prolactin, IGF-II, MIF, and osteopontin) reportedly exhibited an SN of 95.3% at an SP of 99.4% for patients with all stages of ovarian cancer.¹⁹ However, none of the previous studies have evaluated selectivity of panels for ovarian cancer versus benign disease and other malignancies, where selectivity is 1 – SN of the test when evaluated on benign disease and other malignancies for a given SP in controls. Therefore, the need still exists to develop a diagnostic assay that detects stages I and II ovarian cancer with high SN at 98% SP and high selectivity for ovarian cancer in a larger population of patients with early disease. In the present study, we used a multiplexing approach to analyze 96 candidate serum proteins to identify and validate a biomarker combination with the highest power to detect early-stage ovarian cancer and examine its cancer selectivity.

PATIENTS AND METHODS

Patient Populations

The study population was comprised of 2,031 healthy postmenopausal women; 456 patients with ovarian cancer in stage IA ($n = 69$), IB to IIB

($n = 114$), and IIC to IV ($n = 273$); 296 patients with benign pelvic tumors; and postmenopausal women with breast ($n = 210$), lung ($n = 74$), and colorectal ($n = 31$) cancers (Table 1). Samples were obtained from multiple sources (Table 2) and were annotated with information regarding age, cancer diagnosis, stage, histology, and grade (Table 1). The local institutional review boards approved the protocols for use of each sample collection.

Collection and Storage of Blood Serum

Serum samples were collected before surgery and administration of anesthesia. Procedures for serum collection, processing, and storage have been previously described.²⁰ Blood processing was similar for all samples collected at the contributing centers.

Table 2. Sources of Serum Samples

Cancer and Menopausal Status	No. of Samples				
	UPCI	GOG	MDACC	FCCC	Duke UI
Ovarian stage I-IIB, postmenopausal	155	26	2		
Ovarian stage I-II, premenopausal	62				
Ovarian stage IIB-IV, postmenopausal	168	84			21
Benign pelvic, postmenopausal	183	100	13		
Benign pelvic, premenopausal	15				
Breast postmenopausal					210
Lung postmenopausal	37				37
Colorectal postmenopausal	31				
Healthy postmenopausal	46	142	1,783	62	
Healthy premenopausal	63				

Abbreviations: UPCI, University of Pittsburgh Cancer Institute; GOG, Gynecologic Oncology Group Blood Bank; MDACC, The University of Texas M. D. Anderson Cancer Center; FCCC, Fox Chase Cancer Center; Duke, Duke University; UI, University of Iowa.

Sources of Bead-Based Immunoassays

The xMAP bead-based technology (Luminex, Austin, TX) permits multiplexed analysis of several analytes in one sample. Ninety-six bead-based xMAP immunoassays for potential ovarian cancer or epithelial cancer serum biomarkers used in this study are listed in Table 3. All assays were research grade. Antibodies for kallikreins were a generous gift of Eleftherios Diamandis, MD (University of Toronto, Toronto, Ontario, Canada). The inter-assay variability of each assay was 1.5% to 6%. The intra-assay variability for assays performed on the same day was 3% to 9%. The intra-assay variability for assays performed at different days was 5% to 20%, depending on whether the same lot of reagents was used.⁶⁷⁻⁶⁹ Each bead-based assay was validated in comparison with the corresponding enzyme-linked immunosorbent assay based on the same antibody pair and has demonstrated 89% to 98% correlation (data presented on the University of Pittsburgh Web site⁷⁰ for in-house assays; performance of purchased assays was in agreement with that claimed by a manufacturer).

Multiplex Analysis

Assays were performed according to manufacturers' protocols as previously described.²¹ Samples were analyzed using the Bio-Plex suspension array system (Bio-Rad Laboratories, Hercules, CA). For each analyte, 100 beads were analyzed and means were calculated. Analysis of experimental data was performed using four-parameter logistic curve fitting to the standard analyte curves.

Statistical Analysis

All development of statistical models for distinguishing ovarian cancer patient cases from controls was restricted to the training set until one panel and one model of combining the candidate biomarkers in the panel were selected. A Metropolis algorithm with Monte Carlo simulation (MMC) was used for analysis of the data.⁷¹⁻⁷³ In MMC analysis, the scoring function (SF) for a specific biomarker panel was constructed as a linear combination of logarithms of biomarker concentrations (with minimal detectable concentration added to concentration to reduce the effect of large relative errors near zero concentrations). The Monte Carlo optimization was then used to determine the coefficients in this linear combination that, for the training set, provide the best SN at the desired SP. The cutoff was adjusted at the each iteration of parameter estimates to maintain the desired SP. When a range of cutoffs gave the same SP, we used the average cutoff. The advantage of using the number of misdiagnosed cancers as the optimization criterion is that it does not rely on any assumptions regarding the statistical distribution of the data, and as a result, the optimal linear combination is practically insensitive to the presence of outliers and erroneous data. However, the choice of SF does restrict the combination of biomarkers to linear combinations to provide the optimum separation. Therefore, no patient cases were excluded from this analysis. All possible panels consisting of two, three, and four biomarkers were evaluated for SN at 95% SP in the preliminary training set. For each panel size, the 500 panels with the best SN at 95% SP on the full data set were re-estimated with cross validation. For cross validation, 20% of participants were randomly excluded from the data set, and the rest were used as at training set to build the optimal SF. The resultant model was applied to the excluded participants, and this process was repeated 400 times. A smaller subset of biomarkers that comprised the top 20 panels was then re-evaluated in a larger training set at 98% SP. One optimal panel was chosen that offered high cross-validated SN for both early- and late-stage ovarian cancer at 98% SP. This sole panel and method of combination was evaluated in the validation set.

Univariate comparisons were made using the Wilcoxon rank sum test or *t* test for log-transformed data. Exact CIs for binomial proportions are provided by the R function `binom.test`, which inverts the exact binomial test. Comparisons between different samples sources were performed using one-way analysis of variance (ANOVA) with Tukey's multiple comparison test.

Table 3. Multiplexed Biomarkers

Biologic Group and Source*	Proteins	Plex No.†
Cytokines/chemokines		
Invitrogen/Biosource	Eotaxin-1, TNFR1, TNFR2, IL-1R α , IL-2R, IL-6R	1
Millipore/Linco	IL-1b, IL-4, IL-5, IL-6, IL-7, IL-8, IL-10, IL-12p40, IL-13, G-CSF, GM-CSF, IFN γ , TNF- α , IP-10, MCP-1, MIP-1 α , MIP-1 β , MIF, CD40L, fractalkine ^{21,22}	2
Growth/angiogenic/factors		
Invitrogen/Biosource	EGF, ²¹ VEGF, bFGF, HGF, NGF, ²³ EGFR ^{13,24}	1
Luminex Core Facility, University of Pittsburgh	Her2/neu, ^{25,26} IGFBP-1 ²⁷	3
Antiangiogenic factors		
Luminex Core Facility, University of Pittsburgh	Angiostatin, endostatin, thrombospondin ^{28,29}	4
Cancer antigens		
Luminex Core Facility, University of Pittsburgh	CA-125, ^{30,31} CA 15-3, ^{32,33} CEA ³⁴⁻³⁶	5
Luminex Core Facility, University of Pittsburgh	CA 19-9, ³⁷ CA 72-4,37,38 AFP, ³⁹ SCC ⁴⁰	6
Apoptotic proteins		
Luminex Core Facility, University of Pittsburgh	Cyfra 21-1 ^{15,41}	5
Invitrogen/Biosource	DR5 ⁴²	1
Millipore/Linco	Fas, FasL ⁴³	7
Proteases/binding proteins		
Luminex Core Facility, University of Pittsburgh	Kallikreins 8 and 10 ^{44,45}	5, 6
R&D Systems	MMP-1 to MMP-13 ⁴⁶⁻⁴⁸	8
R&D Systems	TIMP-1 to TIMP-4	9
Adhesion molecules		
Millipore/Linco	ICAM, VCAM, E-selectin ⁴⁹	10
Hormones		
Millipore/Linco	Prolactin, TSH, LH, ACTH, GH ^{50,51}	11
Luminex Core Facility, University of Pittsburgh	β HCG ^{52,53}	6
Adipokines		
Millipore/Linco	Adiponectin, leptin, resistin ⁵⁴⁻⁵⁷	12
Other markers		
Luminex Core Facility, University of Pittsburgh	Mesothelin ⁵⁸	6
Luminex Core Facility, University of Pittsburgh	HE4 ⁵⁹	13
Millipore/Linco	tPAI-1, active PAI-1, ⁶⁰ MPO ⁶¹	14
Millipore/Linco	Apolipoproteins (Apo) A1, AII, B, CII, CIII, E ^{18,62,63}	15
Luminex Core Facility, University of Pittsburgh	TTR ¹⁸	16
Millipore/Linco	Insulin, ⁶⁴ osteopontin, ⁶⁶ osteoprotegerin, ⁶⁵ osteocalcin ⁶⁶	17

*Invitrogen/Biosource, Camarillo, CA; Millipore/Linco, St Louis, MO; Luminex Core Facility, University of Pittsburgh, Pittsburgh, PA; and R&D Systems, Minneapolis, MN.

†Plex No. indicates multiplexed panel (i.e., biomarkers that were analyzed simultaneously).

RESULTS

Bead-Based Immunoassay Analysis of Biomarker Serum Concentrations in Patients With Ovarian Cancer

Because of the low prevalence of ovarian cancer, the required minimum SP of serum test should be 98% when combined with TVS




Fig 1. Flowchart of experimental design. MMC, Metropolis algorithm with Monte Carlo simulation; SP, specificity.

as a second-line test.^{10,11} Accurate assessment of such high SP requires at least 1,000 healthy participants. Therefore, our strategy was to first analyze the preliminary smaller training set using MMC algorithm to select a subset of biomarker combinations (panels) providing the highest SN at 95% SP and then to use an expanded training set containing more than 1,000 healthy sera to select and validate a panel offering the highest SN at 98% SP (Fig 1).

Serum samples from the preliminary training set of postmenopausal women consisting of 48 patients with stage IA, 91 patients with stage IB to IIB, and 149 patients with stage III to IV ovarian cancer and 276 age-matched healthy controls were analyzed using multiplexed bead-based immunoassays for 96 cancer-associated biomarkers representing proteins with different biologic functions (Table 3). These candidate biomarkers were selected based on the published evidence documenting an association with epithelial cancer development and progression.^{13,15,18,21-63,74} Serum concentrations of 35 proteins demonstrated highly significant ($P < .001$) differences between ovarian cancer samples and samples from healthy controls (Data Supplement Fig 1). Circulating concentrations of CA-125, HE4, prolactin, IL-2R, CA 72-4, CA 15-3, CA 19-9, MIF, Cyfra 21-1, TNFR1, TNFR2, IL-7, IL-10, IGFBP1, IL-6, TNF- α , GH, TSH, ACTH, TIMP-1, and osteopontin were significantly ($P < .001$) higher in serum of patients with early-stage ovarian cancer compared with healthy women, whereas concentrations of eotaxin-1, MMP-2, MMP-3, VCAM-1, EGFR, ErbB2, FSH, LH, CD40L, ApoA1, adiponectin, TTR, and osteocalcin were significantly ($P < .001$) lower in serum of patients with early-stage ovarian cancer compared with healthy women. Serum levels of HE4, prolactin, IL-2R, CA 72-4, CA 15-3, CA 19-9, Cyfra 21-1, TNFR1, TNFR2, IL-7, IL-10, IGFBP1, IL-6, TNF- α , TSH, MMP-7, eotaxin-1, VCAM-1, ErbB2, FSH, LH, ApoA1, adiponectin, TTR, and CD40L differed significantly ($P < .01$) between patients with early-stage (stages I and II) and late-stage (stages III and IV) ovarian cancer (Data Supplement Fig 1). Circulating concentrations of CA-125, HE4, IL-2R, CA 72-4, CA 15-3, CA 19-9, Cyfra 21-1, TNFR1, TNFR2, IL-7, IL-10, IGFBP1, IL-6, TIMP-1, osteopontin, MMP-7, MMP-2, EGFR, FSH, LH, ApoA1, and TTR were significantly ($P < .05$) different between early ovarian cancer and benign pelvic disease. No significant differences in serum concentrations of any of the biomarkers between stages I and II were found (data not shown).

Statistical Analysis of Multimarker Panels

MMC algorithm was applied to the preliminary training set consisting of sera from patients with ovarian cancer (stage IA, $n = 54$; stage IB to II, $n = 85$; and stage IIC to IV, $n = 149$) and 276 healthy women. Performance of two-, three-, four-, and five-biomarker panels was compared. Four-biomarker panels offered superior performance compared with two- and three-biomarkers panels, whereas using five-biomarker panels did not result in significant improvement. Thirty-five four-biomarker panels with statistically indistinguishable classification accuracy for early-stage disease were identified (Table 4). These panels represented various combinations of the following 16 biomarkers: eotaxin-1, IL-2R, MMP-2, MMP-3, Cyfra 21-1, ErbB2, EGFR, CEA, CA72-4, IGFBP-1, VCAM-1, FSH, GH, HE4, MMP-7, and CA-125. Distributions of 16 biomarkers in samples obtained from different cancer centers were analyzed by one-way ANOVA with Tukey's multiple comparison test to assess bias between the centers (Data Supplement Fig 2). For VCAM-1, eotaxin-1, Cyfra 21-1, ErbB2, EGFR, MMP-3, IGFBP-1, IL-2R, and GH, the differences

Table 4. Sensitivities of Different Panels After Cross Validation at 95% Specificity Using the MMC Algorithm

Marker				Sensitivity (%)
M1	M2	M3	M4	
CA-125	Eotaxin-1	CEA	HE4	96.1
CA-125	CEA	VCAM-1	HE4	95.7
CA-125	Eotaxin-1	EGFR	HE4	94.6
CA-125	Eotaxin-1	IGFBP-1	HE4	94.7
CA-125	Eotaxin-1	HE4	MMP-7	94.2
CA-125	VCAM-1	GH	MMP-7	93.6
CA-125	CA 72-4	VCAM-1	HE4	94.2
CA-125	CEA	IGFBP-1	HE4	94.1
CA-125	Eotaxin-1	ErbB2	HE4	94.3
CA-125	IGFBP-1	VCAM-1	MMP-7	94
CA-125	IGFBP-1	VCAM-1	HE4	94.5
CA-125	Eotaxin-1	VCAM-1	HE4	94.6
CA-125	MMP-3	VCAM-1	MMP-7	93.5
CA-125	EGFR	CEA	HE4	93.7
CA-125	CEA	IL-2R	HE4	94.1
CA-125	Eotaxin-1	IL-2R	HE4	94
CA-125	Eotaxin-1	GH	HE4	94.3
CA-125	VCAM-1	GH	HE4	93.9
CA-125	ErbB2	CEA	HE4	93.6
CA-125	Eotaxin-1	CA 72-4	HE4	94.1
CA-125	Eotaxin-1	IGFBP-1	VCAM-1	93.6
CA-125	Eotaxin-1	CEA	IGFBP-1	93.2
CA-125	Eotaxin-1	MMP-2	HE4	94
CA-125	MMP-2	CEA	HE4	93.9
CA-125	Eotaxin-1	VCAM-1	MMP-7	93.3
CA-125	CEA	CA 72-4	HE4	93.3
CA-125	EGFR	VCAM-1	HE4	94
CA-125	MMP-3	CEA	HE4	93.3
CA-125	Cyfra 21-1	CEA	HE4	93
CA-125	Eotaxin-1	MMP-3	MMP-7	93.2
CA-125	Eotaxin-1	IGFBP-1	GH	92.6
CA-125	MMP-2	VCAM-1	MMP-7	93.2
CA-125	Eotaxin-1	IGFBP-1	MMP-7	93.3
CA-125	Eotaxin-1	MMP-2	IGFBP-1	93
CA-125	Eotaxin-1	IL-2R	HE4	94.1

Abbreviation: MMC, Metropolis algorithm with Monte Carlo simulation.

between sites were not significant, whereas for CA-125, HE4, CEA, MMP-7, FSH, and CA 72-4, statistical significance between samples sources was observed. Even when the differences between sites using values from control participants were statistically significant, the magnitude of the biases was small compared with the differences between cancer patients and controls. For example, for CA-125, the ANOVA was significant, but the point estimates for the differences between sites ranged from 3% to 25%. In comparison, the differences between the cancer patients and controls ranged from 88% to 93%, far larger than the differences between sites.

These biomarkers were reanalyzed in an expanded training set containing 826 additional healthy controls (total $N = 1,102$) to ascertain performance at 98% SP. A panel selected for best performance was CA-125, HE4, CEA, and VCAM-1; this panel provided 86% SN (95% CI, 79% to 91%) for early-stage ovarian cancer and 93% SN (95% CI, 88% to 97%) for late-stage ovarian cancer at 98% SP (Fig 2). The coefficients of the logarithm of the biomarker concentrations were as follows: CA-125, 0.86; HE4, 0.18; CEA, -0.15; and VCAM-1, -1.44. For comparison, the SN of CA-125 alone at 98% SP was 61% (95% CI, 53% to 69%) for early-stage disease and 83% (95% CI, 76% to 89%) for late-stage disease (Fig 3). The four-biomarker panel offered comparable classification for the four most prevalent histologic types of ovarian cancer—serous (SN = 89%), mucinous (SN = 85%), endometrioid (SN = 90%), and clear cell (SN = 85%) carcinoma. Nonepithelial histologies were classified with an SN of 79%.

Next, the classification results were validated in an independent blinded validation set consisting of 929 healthy controls, 296 benign participants, and 168 patients with ovarian cancer (44 patients with stage I to IIB ovarian cancer, including 14 patients with stage IA and 124 patients with stages IIC to IV ovarian cancer; Table 1). The blinding of validation set was performed by the personnel at the sites providing the biospecimens, and the patient case/control status was unblinded only after the algorithm and its predictions for the validation set were provided back to the sites. The optimal four-biomarker panel classified early-stage cancers with 86% SN (95% CI, 73% to 95%) and late-stage cancers with 95% SN (95% CI, 90% to 98%) at 98% SP, which was significantly higher than CA-125 alone. Of 14 patients with stage IA disease, three were misclassified, resulting in 79% SN (95% CI, 49% to 95%). The four-biomarker set performed well for classification of patients with stage III to IV disease, offering 94% SN (95% CI, 79% to 98%) at 98% SP. Figure 2B demonstrates distributions of scores obtained for analysis of different cancer groups and healthy controls in both training and validation sets by MMC algorithm. This multimarker panel correctly diagnosed 67% (95% CI, 61% to 72%) of 296 blinded benign samples as noncancers.

Selectivity of the Panel for Ovarian Cancer versus Other Epithelial Cancers

The panel of CA-125, HE4, CEA, and VCAM-1 was used to classify a blinded mixed set of patient cases with three other common female cancers, breast ($n = 210$), colorectal ($n = 31$), and lung ($n = 74$). This panel correctly identified 94% (95% CI, 90% to 97%) of breast cancers, 100% (95% CI, 89% to 100%) of colorectal cancers, and 64% (95% CI, 52% to 74%) of lung cancers as non-ovarian cancer.




Fig 2. Cumulative receiver operating characteristic (ROC) curves in the training set and distribution of scores in training and validation sets using the four-biomarker panel with Metropolis algorithm with Monte Carlo simulation (MMC): (A) healthy controls versus stage I to IIB ovarian cancer; (B) healthy controls versus stages IIC to IV ovarian cancer. Solid line indicates four-biomarker panel; dashed line indicates CA125. (C) Distributions of scores created by MMC algorithm. P value over a group denotes statistical significance of differences between each group member and appropriate healthy control. Breast, colon, and lung cancers are compared with early-stage ovarian cancer in the validation set.

DISCUSSION

A multimarker approach seems to hold substantial promise for detection of early-stage ovarian cancers.⁵⁶ The goal of the present

study was to select a panel of biomarkers that would provide high SN and SP for distinguishing early-stage ovarian cancer from healthy controls from a large array of serum proteins with different biologic functions representing various aspects of systemic response to a growing tumor.^{12,21,60,75-88} Availability of a bead-based xMAP multiplexing technology was essential for such analysis because sufficient sera would not be available to perform such a large number of conventional enzyme-linked immunosorbent assays.

After performing a multimarker bead-based immunoassay screening, we observed, in agreement with published evidence, elevated serum levels of IGFBP-1, MIF, HE4, IL-2R, CA19-9, CA-125, Cyfra 21-1, CA 72-4, and prolactin and lower levels of FSH, LH, EGFR, TTR, and ApoA1 in patients with ovarian cancer compared with healthy women.^{9,37,41,86-89} In addition, to the best of our knowledge, we are the first to report significant elevation of serum levels of GH and ACTH and lower concentrations of eotaxin-1, immunodetectable MMP-2 and MMP-3, and VCAM-1 in serum of patients with ovarian cancer. The mechanisms behind lower MMP-2, MMP-3, and VCAM-1 levels in patients with ovarian and other cancers have yet to be determined. We have discussed biologic functions of these differentially expressed proteins and their putative roles in pathways associated with ovarian carcinogenesis in our previous report.⁹⁰

Using an MMC algorithm, we have identified a combination of four serum proteins, CA-125, HE4, CEA, and VCAM-1, that offered 86% SN at 98% SP for early-stage ovarian cancer in the solely evaluated, independent, blinded validation set. Analysis of TTR and ApoA1 on the basis of the discovery by Zhang et al¹⁸ was performed. Using our antibody combinations, neither of these two biomarkers offered additional information compared with the multimarker panels presented in Appendix Table A1 (online only).

The high SN of 86% achieved with our four-marker panel for distinguishing women with early-stage ovarian cancer from healthy individuals at 98% SP surpasses the SNs previously reported (70% to 73%).^{9,37,41,86-89} The fact that the multimarker assay offered high accuracy in a heterogeneous validation set that contained different histologies indicates the likely general utility of the assay for the most common histotypes of ovarian cancer. Of note, in agreement with published evidence, we observed a lower incidence of serous histology in early-stage patients.⁹¹ The panel of biomarkers correctly classified 67% of benign lesions as noncancer. The incidence of benign disease is approximately 2.4 benign cases per each ovarian cancer case.⁹² With 67% of benign lesions recognized as noncancer by the four-biomarker panel, only 0.7 benign lesions will be diagnosed as cancer for each true ovarian cancer. SP would be further improved by the subsequent use of TVS to distinguish malignant from benign pelvic lesions in 2% of women classified as at risk for ovarian cancer, further reducing the number of laparotomies for benign disease.⁹³

The three most commonly diagnosed cancers in women are breast, lung, and colorectal.⁹⁴ The current panel was highly selective for ovarian cancer when compared with breast cancer and moderately selective when compared with lung cancer, suggesting that multimarker assays can be cancer specific. Detection of occasional breast and lung cancers might be facilitated by incorporating mammography and spiral chest computed tomography in the second phase of screening for women with a negative TVS.

The performance of this multimarker panel for detecting ovarian cancer in asymptomatic patients remains to be further determined in longitudinal retrospective studies using the serum banks established by major ongoing prospective trials (eg, Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial; Women's Health Initiative; United Kingdom Collaborative Trial of Ovarian Cancer Screening). In these studies, the CA-125, HE4, CEA, and VCAM-1 panel may be further modified by the addition of other biomarkers to achieve optimal lead time before clinical diagnosis and high SN and SP. SN of the panel for ovarian cancer could be further improved by incorporating longitudinal information similar to the method developed for CA-125.⁹⁵

In summary, data obtained in this study suggest that a four-biomarker panel may detect 86% of patients with early-stage ovarian cancer with an SP of 98%. This panel could be used as the first-line test in a two-step strategy for early detection, similar to the approach used in the current United Kingdom Collaborative Trial of Ovarian Cancer Screening where increasing CA-125 triggers TVS.^{30,96}

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Although all authors completed the disclosure declaration, the following author(s) indicated a financial or other interest that is relevant to the subject matter under consideration in this article. Certain relationships marked with a "U" are those for which no compensation was received; those relationships marked with a "C" were compensated. For a detailed description of the disclosure categories, or for more information about ASCO's conflict of interest policy, please refer to the Author Disclosure Declaration and the Disclosures of Potential Conflicts of Interest section in Information for Contributors.

Employment or Leadership Position: Ian Jacobs, United Kingdom Collaborative Trial of Ovarian Cancer Screening/United Kingdom Familial Ovarian Cancer Screening Study (C), University College London Faculty of Biomedicine (C), Women's Health Specialists (C) **Consultant or Advisory Role:** Ian Jacobs, Becton Dickinson (C); Robert C. Bast Jr, Fujirebio Diagnostics (C), Vermillion (C) **Stock Ownership:** None **Honoraria:** None **Research Funding:** Robert C. Bast Jr, Fujirebio Diagnostics **Expert Testimony:** None **Other Remuneration:** Robert C. Bast Jr, Fujirebio Diagnostics

AUTHOR CONTRIBUTIONS

Conception and design: Steven Skates, Elieser Gorelik, Robert C. Bast Jr, Anna E. Lokshin

Provision of study materials or patients: Francesmary Modugno, Jeffrey Marks, Andrew Godwin, Karen Lu, Robert C. Bast Jr, Anna E. Lokshin

Collection and assembly of data: Zoya Yurkovetsky, Robert C. Bast Jr, Anna E. Lokshin

Data analysis and interpretation: Zoya Yurkovetsky, Steven Skates, Aleksey Lomakin, Brian Nolen, Trenton Pulsipher, Elieser Gorelik, Usha Menon, Donna Badgwell, Robert C. Bast Jr, Anna E. Lokshin

Manuscript writing: Zoya Yurkovetsky, Steven Skates, Aleksey Lomakin, Brian Nolen, Elieser Gorelik, Ian Jacobs, Usha Menon, Robert C. Bast Jr, Anna E. Lokshin

Final approval of manuscript: Zoya Yurkovetsky, Steven Skates, Aleksey Lomakin, Brian Nolen, Elieser Gorelik, Ian Jacobs, Usha Menon, Donna Badgwell, Robert C. Bast Jr, Anna E. Lokshin

REFERENCES

1. Boente MP, Schilder R, Ozols RF: Gynecological cancers. *Cancer Chemother Biol Response Modif* 18:418-434, 1999
2. Baker TR, Piver MS: Etiology, biology, and epidemiology of ovarian cancer. *Semin Surg Oncol* 10:242-248, 1994
3. Holschneider CH, Berek JS: Ovarian cancer: Epidemiology, biology, and prognostic factors. *Semin Surg Oncol* 19:3-10, 2000
4. Jacobs I, Bast RC Jr: The CA 125 tumour-associated antigen: A review of the literature. *Hum Reprod* 4:1-12, 1989
5. van Haaften-Day C, Shen Y, Xu F, et al: OVX1, macrophage-colony stimulating factor, and CA-125-II as tumor markers for epithelial ovarian carcinoma: A critical appraisal. *Cancer* 92:2837-2844, 2001
6. Woolas RP, Xu FJ, Jacobs IJ, et al: Elevation of multiple serum markers in patients with stage I ovarian cancer. *J Natl Cancer Inst* 85:1748-1751, 1993
7. Chi DS, Hoskins WJ: Ovarian cancer controversy: When and how to use available screening methods. *Medscape Womens Health* 1:3, 1996
8. DePriest PD, DeSimone CP: Ultrasound screening for the early detection of ovarian cancer. *J Clin Oncol* 21:194-199, 2003 (suppl 10)
9. MacDonald ND, Rosenthal AN, Jacobs IJ: Screening for ovarian cancer. *Ann Acad Med Singapore* 27:676-682, 1998
10. Menon U, Jacobs IJ: Ovarian cancer screening in the general population. *Curr Opin Obstet Gynecol* 13:61-64, 2001
11. Jacobs IJ, Menon U: Progress and challenges in screening for early detection of ovarian cancer. *Mol Cell Proteomics* 3:355-366, 2004
12. Baron AT, Lafky JM, Boardman CH, et al: Serum sErbB1 and epidermal growth factor levels as tumor biomarkers in women with stage III or IV epithelial ovarian cancer. *Cancer Epidemiol Biomarkers Prev* 8:129-137, 1999
13. Baron AT, Cora EM, Lafky JM, et al: Soluble epidermal growth factor receptor (sEGFR/sErbB1) as a potential risk, screening, and diagnostic serum biomarker of epithelial ovarian cancer. *Cancer Epidemiol Biomarkers Prev* 12:103-113, 2003
14. Mazurek A, Niklinski J, Laudanski T, et al: Clinical tumour markers in ovarian cancer. *Eur J Cancer Prev* 7:23-35, 1998
15. Hasholzner U, Baumgartner L, Stieber P, et al: Significance of the tumour markers CA 125 II, CA 72-4, CASA and CYFRA 21-1 in ovarian carcinoma. *Anticancer Res* 14:2743-2746, 1994
16. Perkins GL, Slater ED, Sanders GK, et al: Serum tumor markers. *Am Fam Physician* 68:1075-1082, 2003
17. Skates SJ, Horick N, Yu Y, et al: Preoperative sensitivity and specificity for early-stage ovarian cancer when combining cancer antigen CA-125II, CA 15-3, CA 72-4, and macrophage colony-stimulating factor using mixtures of multivariate normal distributions. *J Clin Oncol* 22:4059-4066, 2004
18. Zhang Z, Bast RC Jr, Yu Y, et al: Three biomarkers identified from serum proteomic analysis for the detection of early stage ovarian cancer. *Cancer Res* 64:5882-5890, 2004
19. Visintin I, Feng Z, Longton G, et al: Diagnostic markers for early detection of ovarian cancer. *Clin Cancer Res* 14:1065-1072, 2008
20. Yurkovetsky Z, Ta'asan S, Skates S, et al: Development of multimarker panel for early detection of endometrial cancer: High diagnostic power of prolactin. *Gynecol Oncol* 107:58-65, 2007
21. Gorelik E, Landsittel DP, Marrangoni AM, et al: Multiplexed immunobead-based cytokine profiling for early detection of ovarian cancer. *Cancer Epidemiol Biomarkers Prev* 14:981-987, 2005
22. Gallagher NJ, Eliopoulos AG, Agathangelou A, et al: CD40 activation in epithelial ovarian carcinoma cells modulates growth, apoptosis, and cytokine secretion. *Mol Pathol* 55:110-120, 2002
23. Merritt WM, Sood AK: Markers of angiogenesis in ovarian cancer. *Dis Markers* 23:419-431, 2007
24. Maihle NJ, Baron AT, Barrette BA, et al: EGF/ErbB receptor family in ovarian cancer. *Cancer Treat Res* 107:247-258, 2002
25. Cheung TH, Wong YF, Chung TK, et al: Clinical use of serum c-erbB-2 in patients with ovarian masses. *Gynecol Obstet Invest* 48:133-137, 1999
26. Marx D, Fattahi-Meibodi A, Kudelka R, et al: Detection of p105 (c-erbB-2, HER2/neu) serum levels by a new ELISA in patients with ovarian carcinoma. *Anticancer Res* 18:2891-2894, 1998
27. Schoen RE, Weissfeld JL, Kuller LH, et al: Insulin-like growth factor-I and insulin are associated with the presence and advancement of adenomatous polyps. *Gastroenterology* 129:464-475, 2005
28. Paley PJ: Angiogenesis in ovarian cancer: Molecular pathology and therapeutic strategies. *Curr Oncol Rep* 4:165-174, 2002
29. Yokoyama Y, Dhanabal M, Griffioen AW, et al: Synergy between angiostatin and endostatin: Inhibition of ovarian cancer growth. *Cancer Res* 60:2190-2196, 2000
30. Bast RC Jr, Badgwell D, Lu Z, et al: New tumor markers: CA125 and beyond. *Int J Gynecol Cancer* 15:274-281, 2005 (suppl 3)
31. Bast RC Jr, Klug TL, St John E, et al: A radioimmunoassay using a monoclonal antibody to monitor the course of epithelial ovarian cancer. *N Engl J Med* 309:883-887, 1983
32. Jacobs IJ, Rivera H, Oram DH, et al: Differential diagnosis of ovarian cancer with tumour markers CA 125, CA 15-3 and TAG 72.3. *Br J Obstet Gynaecol* 100:1120-1124, 1993
33. Scambia G, Benedetti Panici P, Baiocchi G, et al: CA 15-3 as a tumor marker in gynecological malignancies. *Gynecol Oncol* 30:265-273, 1988
34. Buamah PK, Rake MO, Drake SR, et al: Serum CA 12-5 concentrations and CA 12-5/CEA ratios in patients with epithelial ovarian cancer. *J Surg Oncol* 44:97-99, 1990
35. de Bruijn HW, ten Hoor KA, Boonstra H, et al: Cancer-associated antigen CA 195 in patients with mucinous ovarian tumours: A comparative analysis with CEA, TATI and CA 125 in serum specimens and cyst fluids. *Tumour Biol* 14:105-115, 1993
36. Yabushita H, Masuda T, Ogawa A, et al: Combination assay of CA125, TPA, IAP, CEA, and ferritin in serum for ovarian cancer. *Gynecol Oncol* 29:66-75, 1988
37. Gadducci A, Cosio S, Carpi A, et al: Serum tumor markers in the management of ovarian, endometrial and cervical cancer. *Biomed Pharmacother* 58:24-38, 2004
38. Guadagni F, Roselli M, Cosimelli M, et al: CA 72-4 serum marker: A new tool in the management of carcinoma patients. *Cancer Invest* 13:227-238, 1995
39. Seppälä M, Pihko H, Ruoslahti E: Carcinoembryonic antigen and alpha fetoprotein in malignant tumors of the female genital tract. *Cancer* 35:1377-1381, 1975
40. Yamada T, Ikeda A, Okamoto Y, et al: High serum SCC antigen levels in two patients with ovarian cancer. *Acta Obstet Gynecol Scand* 76:283-285, 1997
41. Gadducci A, Ferdeghini M, Cosio S, et al: The clinical relevance of serum CYFRA 21-1 assay in patients with ovarian cancer. *Int J Gynecol Cancer* 11:277-282, 2001
42. Ouellet V, Le Page C, Madore J, et al: An apoptotic molecular network identified by microarray: On the TRAIL to new insights in epithelial ovarian cancer. *Cancer* 110:297-308, 2007
43. Arts HJ, de Jong S, Hollema H, et al: Fas and Fas ligand in cyst fluids, serum and tumors of patients with benign and (borderline) malignant ovarian tumors. *Int J Oncol* 26:379-384, 2005
44. Paliouras M, Borgono C, Diamandis EP: Human tissue kallikreins: The cancer biomarker family. *Cancer Lett* 249:61-79, 2007
45. Clements JA, Willemsen NM, Myers SA, et al: The tissue kallikrein family of serine proteases: Functional roles in human disease and potential as clinical biomarkers. *Crit Rev Clin Lab Sci* 41:265-312, 2004
46. Acar A, Onan A, Coskun U, et al: Clinical significance of serum MMP-2 and MMP-7 in patients with ovarian cancer. *Med Oncol* 25:279-283, 2008
47. Dolo V, D'Ascenzo S, Violini S, et al: Matrix-degrading proteinases are shed in membrane vesicles by ovarian cancer cells in vivo and in vitro. *Clin Exp Metastasis* 17:131-140, 1999
48. Rauvala M, Puistola U, Turpeenniemi-Hujanen T: Gelatinases and their tissue inhibitors in ovarian tumors; TIMP-1 is a predictive as well as a prognostic factor. *Gynecol Oncol* 99:656-663, 2005
49. Banks RE, Gearing AJ, Hemingway IK, et al: Circulating intercellular adhesion molecule-1 (ICAM-1), E-selectin and vascular cell adhesion molecule-1 (VCAM-1) in human malignancies. *Br J Cancer* 68:122-124, 1993
50. Helzlsouer KJ, Alberg AJ, Gordon GB, et al: Serum gonadotropins and steroid hormones and the development of ovarian cancer. *JAMA* 274:1926-1930, 1995
51. Ylikorkala O, Kaupilla A, Rajala T: Pituitary gonadotrophins and prolactin in patients with endometrial cancer, fibroids or ovarian tumours. *Br J Obstet Gynaecol* 86:901-904, 1979
52. Ind T, Iles R, Shepherd J, et al: Serum concentrations of cancer antigen 125, placental alkaline phosphatase, cancer-associated serum antigen and free beta human chorionic gonadotrophin as prognostic markers for epithelial ovarian cancer. *Br J Obstet Gynaecol* 104:1024-1029, 1997
53. Okamoto T, Niu R, Matsuo K, et al: Human chorionic gonadotropin beta-core fragment is directly produced by cancer cells. *Life Sci* 68:861-872, 2001
54. Erturk E, Tuncel E: Polycystic ovarian disease and serum leptin levels? *Fertil Steril* 80:1068-1069, 2003
55. Sieminska L, Marek B, Kos-Kudla B, et al: Serum adiponectin in women with polycystic ovarian syndrome and its relation to clinical, metabolic and endocrine parameters. *J Endocrinol Invest* 27:528-534, 2004
56. Mor G, Visintin I, Lai Y, et al: Serum protein markers for early detection of ovarian cancer. *Proc Natl Acad Sci U S A* 102:7677-7682, 2005
57. Seow KM, Juan CC, Hsu YP, et al: Serum and follicular resistin levels in women with polycystic

ovarian syndrome during IVF-stimulated cycles. *Hum Reprod* 20:117-121, 2005

58. Huang CY, Cheng WF, Lee CN, et al: Serum mesothelin in epithelial ovarian carcinoma: A new screening marker and prognostic factor. *Anticancer Res* 26:4721-4728, 2006

59. Hellström I, Raycraft J, Hayden-Ledbetter M, et al: The HE4 (WFDC2) protein is a biomarker for ovarian carcinoma. *Cancer Res* 63:3695-3700, 2003

60. Abendstein B, Daxenbichler G, Windbichler G, et al: Predictive value of uPA, PAI-1, HER-2 and VEGF in the serum of ovarian cancer patients. *Anticancer Res* 20:569-572, 2000

61. Koziol-Montewka M, Magrys A, Paluch-Oles J, et al: MPO and cytokines in the serum of cancer patients in the context of Candida colonization and infection. *Immunol Invest* 35:167-179, 2006

62. Gadosmska H, Grzechocinska B, Janecki J, et al: Serum lipids concentration in women with benign and malignant ovarian tumours. *Eur J Obstet Gynecol Reprod Biol* 120:87-90, 2005

63. Moore LE, Fung ET, McGuire M, et al: Evaluation of apolipoprotein A1 and posttranslationally modified forms of transthyretin as biomarkers for ovarian cancer detection in an independent study population. *Cancer Epidemiol Biomarkers Prev* 15:1641-1646, 2006

64. Yam D, Ben-Hur H, Fink A, et al: Insulin and glucose status, tissue and plasma lipids in patients with tumours of the ovary or endometrium: Possible dietary implications. *Br J Cancer* 70:1186-1187, 1994

65. Holen I, Shipman CM: Role of osteoprotegerin (OPG) in cancer. *Clin Sci (Lond)* 110:279-291, 2006

66. Pietschmann P, Zielinski C, Woloszczuk W: Serum osteocalcin levels in breast cancer patients. *J Cancer Res Clin Oncol* 115:456-458, 1989

67. Gu Y, Zeleniuch-Jacquotte A, Linkov F, et al: Reproducibility of serum cytokines and growth factors. *Cytokine* 45:44-49, 2009

68. Arslan AA, Gu Y, Zeleniuch-Jacquotte A, et al: Reproducibility of serum pituitary hormones in women. *Cancer Epidemiol Biomarkers Prev* 17:1880-1883, 2008

69. Linkov F, Gu Y, Arslan AA, et al: Reliability of tumor markers, chemokines, and metastasis-related molecules in serum. *Eur Cytokine Netw* 20:21-26, 2009

70. University of Pittsburgh Cancer Institute: Overview. <http://www.upci.upmc.edu/facilities/luminex/index.html>

71. Metropolis N, Rosenbluth AWW, Rosenbluth MN, et al: Equation of state calculations by fast computing machines. *J Chem Phys* 21:1087, 1953

72. Robert CP, Casella G: Monte Carlo Statistical Methods. New York, NY, Springer Verlag, 2004

73. Fishman GS: Monte Carlo: Concepts, Algorithms, and Applications. New York, NY, Springer Verlag, 1995

74. Kucera E, Kainz C, Tempfer C, et al: Prostate specific antigen (PSA) in breast and ovarian cancer. *Anticancer Res* 17:4735-4737, 1997

75. Callet N, Delaunay J, Pichon MF: Serum sICAM-1 (soluble intercellular adhesion molecule-1) and M-CSF (macrophage colony-stimulating growth factor) throughout monitoring of 34 non-serous ovarian cancers. *Eur J Gynaecol Oncol* 21:135-140, 2000

76. Thierry van Dessel HJ, Lee PD, Faessen G, et al: Elevated serum levels of free insulin-like growth factor I in polycystic ovary syndrome. *J Clin Endocrinol Metab* 84:3030-3035, 1999

77. Fahlbusch B, Metzner G, Schumann I, et al: Inhibition of macrophage migration by a factor from ascites fluids of ovarian cancer patients: II. Production and characterization of an anti-MIF antibody. *Biomed Biochim Acta* 45:371-384, 1986

78. Schummer M, Ng WV, Bumgarner RE, et al: Comparative hybridization of an array of 21,500 ovarian cDNAs for the discovery of genes overexpressed in ovarian carcinomas. *Gene* 238:375-385, 1999

79. Borgoño CA, Fracchioli S, Yousef GM, et al: Favorable prognostic value of tissue human kallikrein 11 (hK11) in patients with ovarian carcinoma. *Int J Cancer* 106:605-610, 2003

80. Diamandis EP, Scorilas A, Fracchioli S, et al: Human kallikrein 6 (hK6): A new potential serum biomarker for diagnosis and prognosis of ovarian carcinoma. *J Clin Oncol* 21:1035-1043, 2003

81. Kim H, Scorilas A, Katsaros D, et al: Human kallikrein gene 5 (KLK5) expression is an indicator of poor prognosis in ovarian cancer. *Br J Cancer* 84:643-650, 2001

82. Kishi T, Grass L, Soosaipillai A, et al: Human kallikrein 8, a novel biomarker for ovarian carcinoma. *Cancer Res* 63:2771-2774, 2003

83. Kyriakopoulou LG, Yousef GM, Scorilas A, et al: Prognostic value of quantitatively assessed KLK7

expression in ovarian cancer. *Clin Biochem* 36:135-143, 2003

84. Obiezu CV, Scorilas A, Katsaros D, et al: Higher human kallikrein gene 4 (KLK4) expression indicates poor prognosis of ovarian cancer patients. *Clin Cancer Res* 7:2380-2386, 2001

85. Yousef GM, Diamandis EP: Expanded human tissue kallikrein family: A novel panel of cancer biomarkers. *Tumour Biol* 23:185-192, 2002

86. Tempfer C, Zeisler H, Sliutz G, et al: Serum evaluation of interleukin 6 in ovarian cancer patients. *Gynecol Oncol* 66:27-30, 1997

87. Macciò A, Lai P, Santona MC, et al: High serum levels of soluble IL-2 receptor, cytokines, and C reactive protein correlate with impairment of T cell response in patients with advanced epithelial ovarian cancer. *Gynecol Oncol* 69:248-252, 1998

88. Bonfrer JM, Baan AW, Jansen E, et al: Technical evaluation of three second generation CA 125 assays. *Eur J Clin Chem Clin Biochem* 32:201-207, 1994

89. McIntosh MW, Drescher C, Karlan B, et al: Combining CA 125 and SMR serum markers for diagnosis and early detection of ovarian carcinoma. *Gynecol Oncol* 95:9-15, 2004

90. Nolen B, Marrangoni A, Velikokhatnaya L, et al: A serum based analysis of ovarian epithelial tumorigenesis. *Gynecol Oncol* 112:47-54, 2009

91. Lataifeh I, Marsden DE, Robertson G, et al: Presenting symptoms of epithelial ovarian cancer. *Aust N Z J Obstet Gynaecol* 45:211-214, 2005

92. Westhoff C, Clark CJ: Benign ovarian cysts in England and Wales and in the United States. *Br J Obstet Gynaecol* 99:329-332, 1992

93. Jacobs IJ, Skates SJ, MacDonald N, et al: Screening for ovarian cancer: A pilot randomised controlled trial. *Lancet* 353:1207-1210, 1999

94. Jemal A, Siegel R, Ward E, et al: Cancer statistics, 2008. *CA Cancer J Clin* 58:71-96, 2008

95. Skates SP, Pauler DK, Jacobs IJ: Screening based on the risk of cancer calculation from Bayesian hierarchical change-point and mixture models of longitudinal markers. *J Am Stat Assoc* 96:429-439, 2001

96. Menon U, Skates SJ, Lewis S, et al: Prospective study using the risk of ovarian cancer algorithm to screen for ovarian cancer. *J Clin Oncol* 23:7919-7926, 2005